


Abstract

A compound of formula (I)



R¹ is hydrogen or alkyl optionally substituted by hydroxy, alkoxy, or alkylthio, R² is hydrogen, alkyl, hydroxyalkyl, alkylcarbonyloxyalkyl, alkoxyalkyl, alkylthioalkyl, alkenyl, cycloalkylalkyl, heterocyclalkyl, aralkyl in which the aryl ring thereof is optionally fused to a 5-membered heterocyclic group or is optionally substituted by one or more substituents selected from alkoxy, amino, alkylamino, dialkylamino, acylamino, halogen, hydroxy, aminosulfonyl, alkylaminosulfonyl, dialkylaminosulfonyl, alkylsulfonylamino or dialkylaminosulfonylamino, R³ is hydrogen or alkyl optionally substituted by hydroxy, alkoxy, or alkylthio, R⁴ is hydrogen or alkyl, R⁵ is a quinolinyl, isoquinolinyl or oxodihydroisoquinolinyl group optionally fused to a 5-membered heterocyclic group and optionally substituted by one or more substituents selected from halogen, cyano, hydroxy, alkyl, hydroxyalkyl, alkoxyalkyl, alkylthioalkyl, alkoxy, alkylthio, alkenyl, alkoxycarbonyl, alkynyl, carboxyl, acyl, a group of formula -N(R⁶)R⁷, aryl optionally substituted by one or more substituents selected from halogen or alkoxy, or heteroaryl having 5 or 6 ring atoms attached through a ring carbon atom to the indicated carbon atom, and R⁶ and R⁷ are each independently hydrogen or alkyl optionally substituted by hydroxy or alkoxy or one of R⁶ and R⁷ is hydrogen and the other is acyl, or R⁶ and R⁷ together with the nitrogen atom to which they are attached denote a 5- or 6- membered heterocyclalkyl group.